



ELSEVIER

Discrete Mathematics 171 (1997) 299–300

DISCRETE
MATHEMATICS

Author index to volume 171 (1997)

- Aharoni, R., G.T. Herman and A. Kuba, Binary vectors partially determined by linear equation systems (1–3) 1– 16
- Armanious, M.H., Construction of nilpotent sloops of class n (1–3) 17– 25
- Brandstädt, A., F. F. Dragan and F. Nicolai, LexBFS-orderings and powers of chordal graphs (1–3) 27– 42
- Broersma, H., H. Li, J. Li, F. Tian and H.J. Veldman, Cycles through subsets with large degree sums (1–3) 43– 54
- Bryant, D.E., A. Rodger and E.R. Spicer, Embeddings of m -cycle systems and incomplete m -cycle systems: $m \leq 14$ (1–3) 55– 75
- Chew, K.H., On Vizing's theorem, Adjacency lemma and fan argument generalized to multigraphs (*Note*) (1–3) 283–286
- Clark, L.H. and D. Haile, Remarks on the size of critical edge-chromatic graphs (*Note*) (1–3) 287–293
- da Silva, I.P.F., Note on inseparability graphs of matroids having exactly one class of orientations (1–3) 77– 87
- Dragan, F.F., see A. Brandstädt (1–3) 27– 42
- Du, D.-Z., see P.-J. Wan (1–3) 261–275
- Dvořák, T., I. Havel and P. Liebl, Euler cycles in the complete graph K_{2m+1} (1–3) 89–102
- Egawa, Y., Contractible cycles in graphs with large minimum degree (1–3) 103–119
- Fon-Der-Flaass, D.G., Arrays of distinct representatives — a very simple NP-complete problem (*Note*) (1–3) 295–298
- Gardner, R., B. Micale, M. Pennisi and R. Zijlstra, Cyclic and rotational hybrid triple systems (1–3) 121–139
- Graham, R.L., see P.-J. Wan (1–3) 261–275
- Haile, D., see L.H. Clark (1–3) 287–293
- Hattingh, J.H. and E. Ungerer, Minus k -subdomination in graphs II (1–3) 141–151
- Havel, I., see T. Dvořák (1–3) 89–102
- Herman, G.T., see R. Aharoni (1–3) 1– 16
- Karoński, M. and T. Łuczak, The number of connected sparsely edged uniform hypergraphs (1–3) 153–167
- Kuba, A., see R. Aharoni (1–3) 1– 16
- Li, H., see H. Broersma (1–3) 43– 54
- Li, J., see H. Broersma (1–3) 43– 54
- Liebl, P., see T. Dvořák (1–3) 89–102
- Liu, R.-Y. and L.-C. Zhao, A new method for proving chromatic uniqueness of graphs (1–3) 169–177
- Łuczak, T., see M. Karoński (1–3) 153–167
- McKee, T.A., Clique neighborhoods and nearly chordal graphs (1–3) 179–189
- Micale, B., see R. Gardner (1–3) 121–139
- Minoux, M., Bideterminants, arborescences and extension of the Matrix-Tree Theorem to semirings (1–3) 191–200
- Mitchem, J. and P. Morriss, On the cost-chromatic number of graphs (1–3) 201–211
- Morriss, P., see J. Mitchem (1–3) 201–211
- Nicolai, F., see A. Brandstädt (1–3) 27– 42
- Pennisi, M., see R. Gardner (1–3) 121–139
- Rodger, C.A., see D.E. Bryant, (1–3) 55– 75
- Rusu, I., Building counterexamples (1–3) 213–227

- Santmyer, J.M., A Stirling like sequence of rational numbers (1-3) 229–235
- Spicer, E.R., see D.E. Bryant, (1-3) 55– 75
- Tardif, C., A fixed box theorem for the cartesian product of graphs and metric spaces (1-3) 237–248
- Teschner, U., New results about the bondage number of a graph (1-3) 249–259
- Tian, F., see H. Broersma (1-3) 43– 54
- Ungerer, E., see J.H. Hattingh (1-3) 141–151
- Veldman, H.J., see H. Broersma (1-3) 43– 54
- Wan, P.-J., D.-Z. Du and R.L. Graham, The Steiner ratio for the dual normed plane (1-3) 261–275
- Zhao, L.-C., see R.-Y. Liu (1-3) 169–177
- Zijlstra, R., see R. Gardner (1-3) 121–139
- Zörnig, P., On the line graphs of the complete r -partite graphs (1-3) 277–282